

## RESEARCH

## Open Access



# The basic reproductive ratio of Barbour's two-host schistosomiasis model with seasonal fluctuations

Shu-Jing Gao<sup>1,2</sup>, Hua-Hua Cao<sup>2</sup>, Yu-Ying He<sup>2</sup>, Yu-Jiang Liu<sup>2</sup>, Xiang-Yu Zhang<sup>2</sup>, Guo-Jing Yang<sup>3,4</sup> and Xiao-Nong Zhou<sup>1\*</sup>

## Abstract

**Background:** Motivated by the first mathematical model for schistosomiasis proposed by Macdonald and Barbour's classical schistosomiasis model tracking the dynamics of infected human population and infected snail hosts in a community, in our previous study, we incorporated seasonal fluctuations into Barbour's model, but ignored the effect of bovine reservoir host in the transmission of schistosomiasis. Inspired by the findings from our previous work, the model was further improved by integrating two definitive hosts (human and bovine) and seasonal fluctuations, so as to understand the transmission dynamics of schistosomiasis japonica and evaluate the ongoing control measures in Liaonan village, Xingzi County, Jiangxi Province.

**Methods:** The basic reproductive ratio  $R_0$  and its computation formulae were derived by using the operator theory in functional analysis and the monodromy matrix theory. The mathematical methods for global dynamics of periodic systems were used in order to show that  $R_0$  serves as a threshold value that determines whether there was disease outbreak or not. The parameter fitting and the ratio calculation were performed with surveillance data obtained from the village of Liaonan using numerical simulation. Sensitivity analysis was carried out in order to understand the impact of  $R_0$  on seasonal fluctuations and snail host control. The modified basic reproductive ratios were compared with known results to illustrate the infection risk.

**Results:** The Barbour's two-host model with seasonal fluctuations was proposed. The implicit expression of  $R_0$  for the model was given by the spectral radius of next infection operator. The  $R_0$ s for the model ranged between 1.030 and 1.097 from 2003 to 2010 in the village of Liaonan, Xingzi County, China, with 1.097 recorded as the maximum value in 2005 but declined dramatically afterwards. In addition, we proved that the disease goes into extinction when  $R_0$  is less than one and persists when  $R_0$  is greater than one. Comparisons of the different improved models were also made.

**Conclusions:** Based on the mechanism and characteristics of schistosomiasis transmission, Barbour's model was improved by considering seasonality. The implicit formula of  $R_0$  for the model and its calculation were given. Theoretical results showed that  $R_0$  gave a sharp threshold that determines whether the disease dies out or not. Simulations concluded that: (i) ignoring seasonality would overestimate the transmission risk of schistosomiasis, and (ii) molluscicide is an effective control measure to curtail schistosomiasis transmission in Xingzi County when the removal rate of infected snails is small.

**Keywords:** Schistosomiasis japonica, Mathematical model, Seasonal fluctuation, Basic reproductive ratio, Parameter estimation, Barbour's two-host model

\* Correspondence: zhoun1@chinacdc.cn

<sup>1</sup>National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention; Key Laboratory of Parasite and Vector Biology, MOH, WHO Collaborating Center for Tropical Diseases, Shanghai 200025, China  
Full list of author information is available at the end of the article



© The Author(s). 2017 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

## Background

Schistosomiasis japonica, a parasitic disease caused by *Schistosoma japonicum*, has been in existence in the People's Republic of China (P. R. China) for over 2000 years with considerable public-health and economic significance [1, 2]. A large-scale national schistosomiasis control programme was initiated in the mid-1950s [2, 3], when China's population was approximately 600 million. An estimated 11.8 million people were infected with *S. japonicum* [4]. Sustained control efforts have contributed significantly to the dramatic reduction of both transmission intensity and schistosomiasis distribution in China in the past six decades [5–9]. Recent data have shown that progress toward schistosomiasis elimination encountered difficulties and setback due to high re-infection rates, particularly in lake region of the endemic areas [10]. For instance, schistosomiasis re-emerged shortly after the termination of The World Bank Loan Project (WBLP) at the end of 2001 [1, 5, 11]. In order to achieve the goal of schistosomiasis elimination in P. R. China, Chinese government strengthened the national schistosomiasis control programme in 2004. This made schistosomiasis control top priority along with the list of other communicable diseases such as HIV/AIDS, tuberculosis, and hepatitis B in China [4, 12]. Moreover, a revised strategy to effectively control schistosomiasis by using integrated measures in the national control programme has been implemented since 2005 [11].

Over 40 different species of wild and domestic animals have been identified as definitive hosts of *S. japonicum* [13]. Bovines are the major reservoirs for *S. japonicum* in the lake and marshland regions of southern China [14, 15]. A large number of schistosome-infected bovines are distributed in these regions and they excrete large quantities of *S. japonicum* eggs, the majority of which are deposited near or in the lake [14]. The daily faecal output from a water buffalo (~25 kg) has been estimated to be at least 100 times more than that (~250 g) from an individual human [16, 17]. It was reported that the overall prevalence of *S. japonicum* was 9.6 and 7.2% in water buffalo and cattle, respectively, in 1995 [18]. High prevalence of *S. japonicum* infection in bovine reservoir hosts is believed to be the major factor maintaining active transmission in certain areas. A cluster randomized intervention trial which was designed to compare the control (human treatment) and intervention (human and bovine treatment) in some villages concluded that the incidence of human *S. japonicum* infection is reduced with a decline in the infection rates of water buffaloes [19]. In order to remove bovines as a source of infection, several effective measures, including replacing cattle with farm machinery, isolating marshland, and prohibiting grazing in susceptible areas, have been implemented.

Mathematical modelling is a powerful tool to study the transmission dynamics of schistosomiasis [20]. It was first proposed by Macdonald [21]. Following this pioneering work, many mathematical models have been developed, all of which show great potential in aiding our understanding of the interplay of biology, transmission dynamics and control of schistosomiasis [22–27]. In 1996, Macdonald's model was improved by Barbour [28]. It tracks dynamics of both infected human and snails in a community. Barbour's model has played an important role in evaluating possible control strategies [29]. Traditionally, schistosomiasis models assume that all parameters are constant. A real world environment is obviously non-stationary, and should include seasonal variations in snail population. Infection rates vary seasonally due to natural factors (i.e. changes in moisture and temperature) and social factors (i.e. changes in contact rates). Therefore, it is more realistic to assume that the infection rates are periodic rather than constant. In our previous study [30], we constructed the Barbour's single-host model with seasonal fluctuations (BSHSF model) and calculated the basic reproductive ratio to assess the effect of integrated control measures against schistosomiasis in Liaonan village, Xingzi County, Jiangxi Province. However, the impact of the bovine reservoir host on the transmission of schistosomiasis should not be ignored.

This study aims to further modify the Barbour's two-host model with seasonal fluctuations (BTHSF model) and give an implicit expression of the basic reproductive ratio of schistosomiasis and computation method. Also, we will further establish some indexes that predict prevalence variation characteristics of schistosomiasis, and evaluate the prevention and control strategies for schistosomiasis.

## Methods

### Mathematical formulation

The Barbour's single-host model with seasonal fluctuations (BSHSF model) in Gao et al. [30] was given by the following equations:

$$\begin{cases} \frac{dP}{dt} = a(t)\Delta y(1-P) - gP, \\ \frac{dy}{dt} = b(t)\left(\frac{\sum}{\Delta}\right)P(1-y) - \mu y, \end{cases} \quad (1)$$

All variables of Eq. (1) are described in Table 1. The infection rate  $a(t)$  represents the rate at which a single definitive host becomes infected at unit density of infected snails at time  $t$ . The infection rate  $b(t)$  is the rate at which snails become infected at time  $t$ . The two infection rates can be calculated as follows (see [28]):

**Table 1** Interpretation of the model (1)

Parameter	Interpretation
$a(t)$	the rate of incidence for a single definitive host at unit density of infected snails at time $t$
$b(t)$	the rate at which an infected definitive host causes snail infections at time $t$
$g$	the recovery rate for definitive host infections
$\Delta$	the density of snails
$\Sigma$	the density of definitive hosts
$\mu$	per capita removal rate of infected snails

Abbreviations:  $P$ , the prevalence of infection in the definitive host population;  $y$ , the proportion of infected snails

$$a = a_1 a_2 a_3, \quad b = \beta_1 \beta_2 \beta_3 \beta_4,$$

Here,  $\alpha_1$  denotes the rate at which a host has contact with contaminated water per unit time,  $\alpha_2$  denotes the density of cercariae,  $\alpha_3$  denotes the probability that an encounter with cercaria leads to the establishment of an adult parasite,  $\beta_1$  is the rate of egg-laying,  $\beta_2$  is the probability of an egg developing into a miracidium,  $\beta_3$  is the probability that a miracidium penetrates a snail,  $\beta_4$  is the probability that miracidial penetration into an uninfected snail develops into infection.

Seasonality was taken into consideration by assuming that the infection rates  $a$  and  $b$  in model (1) were time-varying and periodic.  $a(t)$ ,  $b(t)$  were taken as the absolute value of sine functions with 365 days period, while the average values of functions  $a(t)$  and  $b(t)$  per year (365 days) were assumed to be equal to the values of parameters  $a$  and  $b$  in [31], respectively.

Furthermore, we considered the impacts of bovine reservoir host on the transmission of schistosomiasis. The BSHSF model (1) was modified and used to create the BTHSF model. The dynamics of the model are governed by the following differential equations:

$$\begin{cases} \frac{dP_1}{dt} = a_1(t)\Delta y(1-p_1) - g_1 P_1, \\ \frac{dP_2}{dt} = a_2(t)\Delta y(1-p_2) - g_2 P_2 \\ \frac{dy}{dt} = \left( \frac{b_1(t)\Sigma_1 P_1 + b_2(t)\Sigma_2 P_2}{\Delta} \right) (1-y) - \mu y. \end{cases} \quad (2)$$

All variables of Eq. (2) are described in Table 2, where  $a_1(t)$  denotes the rate of incidence for one human host at unit density infected by one infectious snail per unit time (one day) at time  $t$ ,  $a_2(t)$  denotes the rate of incidence for one bovine host at unit density infected by one infectious snail per unit time (1 day) at time  $t$ ,  $b_1(t)$  is the rate of incidence for one susceptible snail at unit density infected by one infected human per unit time (1 day) at time  $t$ ,  $b_2(t)$  is the rate of incidence for one susceptible snail at unit density infected by one

**Table 2** Interpretation of the model (2)

Parameter	Interpretation
$a_1(t)$	the infection rate from snail to human at time $t$
$a_2(t)$	the infection rate from snail to bovine at time $t$
$b_1(t)$	the infection rate from human to snail at time $t$
$b_2(t)$	the infection rate from bovine to snail at time $t$
$g_1$	the recovery rate for human host infections
$g_2$	the recovery rate for bovine host infections
$\Delta$	the density of snails
$\Sigma_1$	the density of human hosts
$\Sigma_2$	the density of bovine hosts
$\mu$	per capita removal rate of infected snails

Abbreviations:  $P_1$ , prevalence of infection in the human host population;  $P_2$ , the prevalence of infection in the bovine host population;  $y$ , the proportion of infected snails

infected bovine per unit time (1 day) at time  $t$ . In this study, we assume that  $b_1$  is proportional to  $b_2$ , i.e.  $b_2 = kb_1$ , where  $k$  is the coefficient ratio. According to Zhao et al. [32]:

$$k = \frac{b_2}{b_1} = \frac{A_2 \times B_2}{A_1 \times B_1},$$

where  $A_i$  ( $i=1,2$ ) denote faeces output per human and bovine,  $B_i$  ( $i=1,2$ ) denote the epg of human and bovine, respectively, here epg is the number of eggs per gram of faeces. The number of eggs per bovine per day is more than  $10^5$  [33], and the number of eggs per human per day is about  $\sim 2800$ – $3000$  [34]. Furthermore, many effective measures, such as improved sanitation and health education are implemented to control the spread of schistosomiasis among human beings. In this paper, we choose  $k=50$ . We also take the infection rates as:

$$\begin{cases} a_1(t) = a_{10} |\sin \pi t / 365|, \\ a_2(t) = a_{20} |\sin \pi t / 365|, \\ b_1(t) = b_{10} |\sin \pi t / 365|, \\ b_2(t) = a_{20} |\sin \pi t / 365|. \end{cases}$$

Since infection rates  $a_i$  and  $b_i$  ( $i=1,2$ ) denote the average values of  $a_i(t)$  and  $b_i(t)$  per year, respectively, we have

$$a_1 = \frac{1}{365} \int_0^{365} a_1(t) dt, \quad a_2 = \frac{1}{365} \int_0^{365} a_2(t) dt, \quad \text{and} \quad b_1 = \frac{1}{365} \int_0^{365} b_1(t) dt.$$

Thus we can derive  $a_{10} = \frac{\pi}{2} a_1$ ,  $a_{20} = \frac{\pi}{2} a_2$ ,  $b_{10} = \frac{\pi}{2} b_1$ ,  $b_{20} = \frac{\pi}{2} kb_1$  and

$$\begin{cases} a_1(t) = \frac{\pi}{2} a_1 |\sin \pi t / 365|, \\ a_2(t) = \frac{\pi}{2} a_2 |\sin \pi t / 365|, \\ b_1(t) = \frac{\pi}{2} b_1 |\sin \pi t / 365|, \\ b_2(t) = \frac{\pi}{2} k b_2 |\sin \pi t / 365|. \end{cases}$$

### The basic reproductive ratio ( $R_0$ )

For epidemiological models, the basic reproductive ratio, which is often introduced as a threshold parameter, is defined as the expected number of secondary infections produced by a single infective individual in a completely susceptible population during its entire infectious period [35].  $R_0$  has often been used to predict the trend of the disease transmission and also to assess the effects of control measures. Here, we give the implicit formula  $R_0$  of model (2). It is expected that the infection will persist if  $R_0$  is greater than one. As  $R_0$  increases, the spreading rate of the disease also increases. This implies that more intervention efforts or improved control measures should be implemented. We have also shown that  $R_0$  is a sharp threshold value which determines whether the disease dies out or not in Additional file 1: Appendix B. That is, if  $R_0 < 1$ , then the disease will die out, whereas if  $R_0 > 1$ , the disease will be endemic.

With reference to Bacaër [36], a biologically meaningful threshold value  $R_0$  of BTHSF model (2) was derived by using operator theory in functional analysis and the monodromy matrix of linear periodic system theory. The numerical computation of the basic reproductive ratio was carried out using the mathematical programming language MATLAB 7.1. Also, sensitivity analysis was carried out in order to assess the impact of seasonal fluctuations on  $R_0$ .

### Study area and estimation of model parameters

Liaonan village, Xingzi County, Jiangxi Province was selected as the study area. The basic reproductive ratio was calculated based on the parameters in model (2). Determination of accurate model parameters was the priority when this threshold was applied. The details of the technique used in estimating the parameters have been described in our previous study [30]. In this paper, part of the parameters in model (2) were determined according to their biological significance. In addition, the remaining parameters were estimated based on the annual report surveillance data of Xingzi County from 2003 to 2010. The meanings of parameters  $g_i$ ,  $\Sigma_i$  ( $i = 1, 2$ ),  $\mu$ ,  $\omega$  are consistent with those reported in [30] (see Tables 1 and 2).

The parameters are stated as follows;  $g_1 = 0.00093$  per day,  $\mu = 0.0055$  per day, and  $g_2 = 0.00183$  per day (the data

are given by Wu [31]). The density of snails,  $\Delta$  (/m<sup>2</sup>), and the density of definitive hosts,  $\Sigma_1$ ,  $\Sigma_2$  (/m<sup>2</sup>), were estimated from annual report surveillance data (see Table 3).

It is assumed that the coefficients of model (2) are constant. By setting the right-hand side of equations in model (2) to zero, the steady-state (equilibrium) values for the prevalence of infection in humans, bovines and snails were obtained [28]

$$\frac{1}{1-\bar{y}} = \frac{t_{MS}^{(1)}}{\bar{y} + 1/t_{SM}^{(1)}} + \frac{t_{MS}^{(2)}}{\bar{y} + 1/t_{SM}^{(2)}}, \quad (3)$$

$$\bar{P}_i = \frac{\bar{y}}{\bar{y} + 1/t_{SM}^{(i)}}, \quad (4)$$

where  $t_{MS}^{(i)} = b_i \Sigma_i / (\mu \Delta)$  and  $t_{SM}^{(i)} = a_i \Delta / g_i$  ( $i = 1, 2$ ) are the transmission factors for definitive host  $i$ . By solving (3) and (4), we obtain the prevalence of infection in humans, bovines and snails from 2003 to 2010.

## Results

### Calculation of $R_0$

Following the results of Bacaër [36] and Wang & Zhao [37], we show that  $R_0$  is equal to the value of  $\lambda$ , where  $\lambda$  is the positive root of the equation

$$\rho(W(365, 0, \lambda)) = 1, \quad (5)$$

Here

$$W(365, 0, \lambda) = \exp \left[ \int_0^{365} \begin{pmatrix} -g_1 & 0 \\ 0 & -g_2 \end{pmatrix} dt + \begin{pmatrix} a_1(t)\Delta/\lambda & 0 \\ a_2(t)\Delta/\lambda & -\mu \end{pmatrix} dt \right], \forall \lambda > 0$$

and  $\rho(W(365, 0, \lambda))$  is the spectral radius of matrix  $W(365, 0, \lambda)$ .

The mathematical details that were used to derive the expression for the basic reproductive ratio can be found in Additional file 1: Appendices A, B (Derivation of  $R_0$  and proof of main results).

### Numerical simulation

Based on the annual report data for Xingzi County from 2003 to 2010, the equilibrium values  $(\bar{P}_1, \bar{P}_2, \bar{y})$  (prevalence of infection in human host, bovine reservoir host and snail) are shown in Table 3. In view of the parameters' values given above (see Table 3), the infection rates  $a_1$ ,  $a_2$ ,  $b_1$  and  $b_2$  of BTH model was derived from Eqs. (3) and (4), and are shown in Table 4.

In Additional file 1: Appendix A, we illustrate that  $R_0$  is the positive root of the equation  $\rho(W(365, 0, \lambda)) = 1$ . Using the data in Tables 3 and 4, the values of  $R_0$  for

**Table 3** The data for  $\Delta$ ,  $\Sigma_1$ ,  $\Sigma_2$ ,  $\bar{P}_1$ ,  $\bar{P}_2$  and  $\bar{y}$  from annual report data in the village of Liaonan, Xingzi County, Jiangxi Province, China

Parameter	2003	2004	2005	2006	2007	2008	2009	2010
$\Delta$	27.997	33.9230	91.4602	133.5238	41.0615	17.4960	11.3121	2.4183
$\Sigma_1$	0.0300	0.0199	0.0475	0.0167	0.0309	0.0178	0.0309	0.0176
$\Sigma_2$	0.0013	0.0007	0.0020	0.0006	0.0010	0.0006	0.0013	0.0008
$\bar{P}_1$	0.0671	0.0533	0.0631	0.0454	0.0454	0.0454	0.04530	0.0457
$\bar{P}_2$	0.0283	0.1379	0.1756	0.1144	0.1084	0.0676	0.0639	0.0466
$\bar{y}$	0.0002	0.00005	0.0001	0.0001	0.0002	0.0005	0.0008	0.0015

BTHSF model were calculated and shown in Table 5. Let  $R_0$  be the basic reproductive ratio for BSHSF model. It was calculated using established methods (see Table 6).

Barbour [28], proposed the following model without seasonality (BTH model):

$$\begin{cases} \frac{dP_1}{dt} = a_1\Delta y(1-P_1) - g_1P_1, \\ \frac{dP_2}{dt} = a_2\Delta y(1-P_2) - g_2P_2, \\ \frac{dy}{dt} = \left( \frac{b_1\Sigma_1P_1 + b_2\Sigma_2P_2}{\Delta} \right) (1-y) - \mu y, \end{cases}$$

and obtained the basic reproductive ratio

$$R_0 = \frac{a_1b_1\Sigma_1}{g_1\mu} + \frac{a_2b_2\Sigma_2}{g_2\mu}.$$

Using the data in Tables 3 and 4, we obtain the values of  $R_0$  in the village of Liaonan from 2003 to 2010.

$R_0$  for BTHSF model is shown in Table 3, and it clearly shows that the values range between 1.030 and 1.097 from 2003 to 2010 in the village of Liaonan. Moreover,  $R_0$  for BTH model is always greater than that of BTHSF model at the same time. Furthermore, the variation tendency of  $R_0$  for BTHSF model is shown in Fig. 1.

The prevalence of two hosts and vector host has strong effect on  $R_0$  for BTHSF model. Except in 2003 and 2010, the values of  $R_0$  for BTHSF model are greater than those for BSHSF model in the other years (see Table 4). The prevalence of infected bovines in 2003 and 2010 are less than the other years (see Table 2). This phenomenon illustrates that if the prevalence ratio of infected bovine hosts and infected human hosts ( $\bar{P}_2/\bar{P}_1$ ) is small,  $R_0$  for BTHSF model may be smaller than for BSHSF model. Moreover, we note that

$R_0$  and the prevalence of infection in bovines ( $\bar{P}_2$ ) for BTHSF model are higher in 2004 and 2005 among the years under study as shown in (Tables 1 and 3).

### Sensitivity analysis

In model (2), infection rates  $a_1(t)$  and  $b_1(t)$  ( $i = 1, 2$ ) were taken as sine functions of 365 days period, with the form  $a_1(t) = \frac{\pi}{2}a_1|\sin\pi t/365|$  and  $b_1(t) = \frac{\pi}{2}b_1|\sin\pi t/365|$ . Thus,  $\pi a_i/2$  and  $\pi b_i/2$  reflect the amplitude of infection rates  $a_i(t)$  and  $b_i(t)$ , respectively. We fix  $g_1 = 0.00093$ ,  $g_2 = 0.00183$ ,  $\Delta = 91.4602$ ,  $\Sigma_1 = 0.0475$ ,  $g_1 = 0.00093$ ,  $\Sigma_2 = 0.0020$ ,  $\mu = 0.0055$ ,  $a_2 = 0.0426$ ,  $b_2 = 0.1224$ , and vary  $a_1$  and  $b_1$  in  $[0, 0.04]$  in (5), with other parameters unchanged as above, by numerical simulations we obtain the curve of  $R_0$  with respect to  $a_1$  and  $b_1$  (see Fig. 2). It shows that the larger  $a_1$  and  $b_1$  are the higher  $R_0$  is. When  $a_1$  and  $b_1$  are near to 0,  $R_0$  is less than 1, which indicates that the control strategy is very effective and disease transmission will be interrupted. This implies that  $R_0$  is very sensitive to the changes in  $a_1$  and  $b_1$  when their values are higher than 0.03. Next, if we fix  $a_1 = 0.0068$ ,  $b_1 = 0.0024$ , and let  $a_2$  and  $b_2$  vary. A graph indicating the relationship between  $R_0$  and  $a_2$  and  $b_2$  was obtained (Fig. 3). This graph shows that  $R_0$  increases with an increase in the amplitude of  $a_2$  and  $b_2$ . Finally, as  $\mu$  varies in  $[0, 0.012]$  with other parameters unchanged as above, numerical simulations provide the relationship between  $R_0$  and  $\mu$  (Fig. 4). Figure 4 shows that  $R_0$  is more sensitive to changes in  $\mu$  when  $\mu$  is less than 0.002.

### Discussion

This paper proposed Barbour's two-host model with seasonality (BTHSF model), which is an improvement on

**Table 4** The calculated values of composite parameters  $a_1$ ,  $a_2$ ,  $b_1$  and  $b_2$ 

Parameters	2003	2004	2005	2006	2007	2008	2009	2010
$a_1$	0.0116	0.0309	0.0051	0.0024	0.0061	0.0050	0.0051	0.0120
$a_2$	0.0093	0.1726	0.0320	0.0128	0.0305	0.0151	0.0144	0.0241
$b_1$	0.0082	0.0015	0.0032	0.0260	0.0057	0.0171	0.0085	0.0075
$b_2$	0.4084	0.0754	0.1606	1.2996	0.2871	0.8565	0.4268	0.3725



**Table 5** The result of the basic reproductive ratio from annual report data in the village of Liaonan, Xingzi County, Jiangxi Province, China

The basic reproductive ratios	2003	2004	2005	2006	2007	2008	2009	2010
$R_0$ (BTHSF model)	1.031	1.074	1.097	1.060	1.057	1.038	1.037	1.030
$\mathfrak{R}_0$ (BTH model)	1.052	1.142	1.192	1.114	1.107	1.066	1.064	1.050

the models BTH and BSHSF. Following the idea of Wang & Zhao [37], the implicit formula of  $R_0$  for BTHSF was given. By computation, the annual changing trend of  $R_0$ s in Liaonan village from 2003 to 2010 was given (see Fig. 1). It showed that  $R_0$  peaked in 2005 at 1.09 and declined dramatically until 2010. There are three possible factors responsible for this decline. First, in 2001, the World Bank Loan Project (WBLP) for schistosomiasis control (1992–2001) was terminated. This led to the resurgence of schistosomiasis transmission after 2001 [38, 39]. Secondly, the implementation of integrated control strategies with emphasis on infection source control commenced in 2005. Finally, the high prevalence of *S. japonicum* in bovines contributes to the high value of  $R_0$ .

By numerical simulation, we realize that the basic reproductive number for Barbour's model with constant infection rates is always higher than that of periodic infection rates. This illustrates that ignoring seasonal fluctuations would lead to overestimating schistosomiasis transmission risk.

Chemotherapy, health education, provision of water, sanitation and hygiene, bovine control and snail host control are the main measures for schistosomiasis control. We know that the implementation of chemotherapy strategy is beneficial to the recovery and well-being of patients and cattle. This would result in increasing the parameters  $g_1$  and  $g_2$  (recovery rates) in model (2). Accordingly, the implementation of control strategies, such as health education, provision of water, sanitation and hygiene, bovine control would lead to decrease in; the host's contact rate with contaminated water, the probability that an encounter with cercariae results in the development of an adult parasite and the rate at which schistosome eggs are laid. Therefore, the infection rates ( $a_1(t)$ ,  $a_2(t)$ ,  $b_1(t)$ ,  $b_2(t)$ ) in model (2) would decrease. More so, the implementation of snail control strategy will increase the removal rate of the snail ( $\mu$ ). Due to the implementation of preventive chemotherapy in Xingzi

County, we assessed the sensitivity analysis of the average infection rates and removal rate. Numerical results have shown that  $R_0$  is a monotone increasing function of the four infection rates and a monotone decreasing function of the removal rate. This implies that as the infection rates increases or the removal rate decreases,  $R_0$  increases. Our results also indicate that mollusciciding is an effective means of controlling schistosomiasis transmission in Xingzi County when the removal rate of snails is small.

Finally, in line with the theoretical concept of this paper, we introduce several indexes that were explored extensively. Similar to the methods established in [28],  $R_0$ s are estimated from prevalence data and assumed to be at a steady equilibrium state, so they are always more than 1. In reality,  $R_0$  may be less than 1. However, we can study the transmission intensity by using the values of  $R_0$ . For example, if we define the relative basic reproductive ratio as:

$$R_i^j = \frac{R_{0j}}{R_{0i}}$$

where  $R_{0k}$  denotes the basic reproductive ratio of year  $k$ . If  $R_i^j > 1$ , the transmission status of schistosomiasis in year  $j$  is more serious than year  $i$  and vice versa.


Next, we define the relative mean basic reproductive ratio as

$$\mathfrak{R}_m = \frac{R_{0m}}{\bar{R}_0}$$

Here,  $\bar{R}_0 = \frac{1}{j-i+1} \sum_{k=i}^j R_{0k}$ .  $\mathfrak{R}_m > 1$  shows that the basic reproductive ratio in year  $m$  is greater than the average ratio from year  $i$  to  $j$ , and vice versa. Taking Liaonan village as the example, the relative basic reproductive ratio in 2005 and 2006 is defined as:

**Table 6** The result of the basic reproductive ratio from annual report data in the village of Liaonan, Xingzi County, Jiangxi Province, China

The basic reproductive ratios	2003	2004	2005	2006	2007	2008	2009	2010
$R_0$ (BTHSF model)	1.031	1.074	1.097	1.060	1.057	1.038	1.037	1.030
$R_0$ (BSHSF model)	1.043	1.035	1.040	1.031	1.031	1.031	1.031	1.032



$$R_{2005}^{2006} = \frac{R_{0_{2006}}}{R_{0_{2005}}} = \frac{1.12}{1.20} = 0.9 < 1.$$

This illustrates that the epidemic trend in 2005 is relatively strong, compared to 2006. In addition,



$$\bar{R}_0 = \frac{1}{8} \sum_{k=2003}^{2010} R_{0_k} = 1.11,$$

and

$$\mathfrak{R}_{2010} = \frac{R_{0_{2010}}}{\bar{R}_0} = \frac{1.06}{1.11} = 0.96 < 1.$$

This indicates that the epidemic trends in 2010 are slower than the average level from 2003 to 2010.

Similarly, it is also able to measure the disease transmission situation between two different regions. Let  $R_0^i$ ,  $\tilde{R}_0^i$  denote the basic reproductive ratios of two different




regions in year  $i$ , respectively. The index of the basic reproductive ratio is defined as:

$$\gamma_i = \frac{R_0^i}{\tilde{R}_0^i}.$$

$\gamma_i$  can be used to compare the effect of control and intervention between two regions at the same time. However, many factors such as heterogeneity of definitive host, drug resistance and optimal control method are yet to be considered in this mathematical model. These issues need to be addressed in the future studies.

## Conclusions

We modified and improved Barbour's model to investigate the role of seasonal fluctuations in the spread of



schistosomiasis. The formula for the basic reproductive number  $R_0$  with a single host was generalized with that of two hosts. Theoretical results showed that it is the basic reproductive ratio which is the threshold value that determines whether schistosomiasis should persist or not. The values of  $R_0$  in Liaonan village from 2003 to 2010 were calculated using the mathematical programming language MATLAB 7.1. The results from this study have shown that ignoring seasonality would overestimate the transmission risk of schistosomiasis, and mollusciciding would be the most effective control measure to curtail schistosomiasis transmission in Xingzi County. Also, the indexes established in this paper will lead to the re-evaluation of control strategies between different areas.

## Additional file

**Additional file 1:** Appendix A. Derivation of  $R_0$ . Appendix B. Proof of the main results. (PDF 67 kb)

## Abbreviations

BSH model: Barbour's single-host model; BSHSF model: Barbour's single-host model with seasonal fluctuations; BTH model: Barbour's two-host model; BTHSF model: Barbour's two-host model with seasonal fluctuations; epg: Eggs per gram; ERC: Institutional Ethics Review Committee;  $R_0$ : The basic reproductive ratio; WBLP: The World Bank Loan Project

## Acknowledgments

This study was supported by Dr Bo Tao and his team from the Schistosomiasis Control Station of Xinzhi County, Jiangxi province, People's Republic of China, on the field work and data collection. Thank also to Dr. Eniola M. Abe for his assistance on language corrections.

## Funding

This project was supported by the Natural Science Foundation of China (grant no. 11561004) and the National Natural Science Foundation of China (grant no. 11261004) through SJG, and by the National Natural Science Foundation of China (grant no. 81573261) through GJY, as well as by the National Science and Technology Major Program (grant no. 2012ZX10004 220) and Shanghai Science and Technology Committee (grant no. 11XD1405400) through XNZ.

## Availability of data and materials

The datasets supporting the conclusions of this article are included within the article and its additional file.

## Authors' contributions

SJG and XNZ conceived and designed the study; YJL contributed with numerical simulation and performed the drawing; all authors contributed to the analysis of the data and to writing the manuscript, and gave final approval for publication. All authors read and approved the final manuscript.

## Competing interests

The authors declare that they have no competing interests.

## Consent for publication

Not applicable.

## Ethics approval and consent to participate

This study was approved by the Institutional Ethics Review Committee (ERC) of the National Institute of Parasitic Diseases, China CDC. All the data are from previous databases or yearbooks, and did not include individual data. Therefore, the informed consent was not necessary in this study.

## Author details

<sup>1</sup>National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention; Key Laboratory of Parasite and Vector Biology, MOH, WHO Collaborating Center for Tropical Diseases, Shanghai 200025, China. <sup>2</sup>Key Laboratory of Jiangxi Province for Numerical Simulation and Emulation Techniques, Gannan Normal University, Ganzhou 341000, China. <sup>3</sup>Jiangsu Institute of Parasitic Diseases, Key Laboratory on Control Technology for Parasitic Diseases, Ministry of Health, Wuxi, Jiangsu 214064, China. <sup>4</sup>Swiss Tropical and Public Health Institute, Basel, Switzerland.

Received: 19 April 2016 Accepted: 12 January 2017

Published online: 25 January 2017

## References

1. Zhou XN, Wang LY, Chen MG, Wu XH, Jiang QW, Chen XY, et al. The public health significance and control of schistosomiasis in China - then and now. *Acta Trop.* 2005;96:97–105.
2. Zhang Z, Clark AB, Bivand R, Chen Y, Carpenter T, Peng W, et al. Nonparametric spatial analysis to detect high-risk regions for schistosomiasis in Guichi, China. *Trans R Soc Trop Med Hyg.* 2009;103:1045–52.
3. Chen X, Wang L, Cai J, Zhou XN, Zheng J, Guo J, et al. Schistosomiasis control in China: the impact of a 10-year World Bank Loan Project (1992–2001). *Bull World Health Organ.* 2005;83:43–8.
4. Report Inter-country meeting on strategies to eliminate schistosomiasis from the Eastern Mediterranean Region. Muscat, Oman. 2007.
5. Zhou XN, Guo JG, Wu XH, Jiang QW, Zheng J, Dang H, et al. Epidemiology of schistosomiasis in the People's Republic of China, 2004. *Emerg Infect Dis.* 2007;13:1470–6.
6. Shrivastava J, Qian BZ, Mcvean G, Webster JP. An insight into the genetic variation of *Schistosoma japonicum* in mainland China using DNA microsatellite markers. *Mol Ecol.* 2005;14:839–49.
7. Engels D, Chitsulo L, Montresor A, Savioli L. The global epidemiological situation of schistosomiasis and new approaches to control and research. *Acta Trop.* 2002;82:139–46.
8. Bergquist R, Tanner M. Controlling schistosomiasis in Southeast Asia: a tale of two countries. *Adv Parasitol.* 2010;72:109–44.
9. Zhao QP, Jiang MS, Dong HF, Nie P. Diversification of *Schistosoma japonicum* in mainland China revealed by mitochondrial DNA. *PLoS Neglect Trop Dis.* 2012;6:e1503.
10. Seto EYW, Remais JV, Carlton EJ, Wang S, Liang S, Brindley PJ, Qiu D, et al. Toward sustainable and comprehensive control of schistosomiasis in China: lessons from Sichuan. *PLoS Neglect Trop Dis.* 2011;5:e1372.
11. Li SZ, Luz A, Wang XH, Xu LL, Wang Q, Qian YJ, et al. Schistosomiasis in China: acute infections during 2005–2008. *Chinese Med J Eng.* 2009;122:1009–14.
12. Engels D, Wang LY, Palmer KL. Control of schistosomiasis in China. *Acta Trop.* 2005;96:67–8.
13. Chen MG, Zheng F. Schistosomiasis control in China. *Parasitol Int.* 1999;48:11–9.
14. Guo JG, Li YS, Gray D, Ning A. A drug-based intervention study on the importance of buffaloes for human *Schistosoma japonicum* infection around Poyang Lake, People's Republic of China. *Am J Trop Med Hyg.* 2006;74:335–41.
15. Yang G, Ross AGP, Lin DD, Williams GM, Chen HG, Li Y, et al. A baseline study on the importance of bovines for human *Schistosoma japonicum* infection around Poyang Lake, China. *Am J Trop Med Hyg.* 2001;65:272–8.
16. Ross AGP, Sleight AC, Li Y, Davis GM, Williams GW, Jiang Z, et al. Schistosomiasis in the People's Republic of China: prospects and challenges for the 21st century. *Clin Microbiol Rev.* 2001;14:270–95.
17. He YX, Salafsky B, Ramaswamy K. Host-parasite relationships of *Schistosoma japonicum* in mammalian hosts. *Trends Parasitol.* 2001;17:320–4.
18. Dakui E. Epidemic status of schistosomiasis in China, a nation-wide sampling survey in 1995. Nanjing: Nanjing University Press; 1998.
19. Gray DJ, Williams GM, Li Y, Chen H, Forsyth S, Li RS, et al. A Cluster-randomised intervention trial against *Schistosoma japonicum* in the Peoples' Republic of China: Bovine and human transmission. *PLoS One.* 2009;4:e5900.
20. Webb GF, Browne CJ. A model of the Ebola epidemics in West Africa incorporating age of infection. *J Biol Dynam.* 2016;10:18–30.
21. MacDonald G. The dynamics of helminth infections, with special reference to schistosomes. *Trans R Soc Trop Med Hyg.* 1965;59:489–506.



22. Williams GM, Sleight AC, Li Y, Feng Z, Davis GM, Chen H, et al. Mathematical modelling of schistosomiasis japonica: comparison of control strategies in the People's Republic of China. *Acta Trop.* 2002;82:253–62.
23. Feng Z, Li CC, Milner FA. Schistosomiasis models with two migrating human groups. *Math Comput Model.* 2005;41:1213–30.
24. Chen ZM, Zou L, Shen DW, Zhang WN, Ruan SG. Mathematical modelling and control of schistosomiasis in Hubei Province, China. *Acta Trop.* 2010; 115:119–25.
25. Zhang P, Feng Z, Milner F. A schistosomiasis model with an age-structure in human hosts and its application to treatment strategies. *Math Biosci.* 2007; 205:83–107.
26. Das P, Mukherjee D, Sarkar AK. A study of schistosome transmission dynamics and its control. *J Biol Syst.* 2006;14:295–302.
27. Gray DJ, Williams GM, Li YS, McManus DP. Transmission dynamics of *Schistosoma japonicum* in the lakes and marshlands of China. *PLoS One.* 2008;3:e4058.
28. Barbour AD. Modeling the transmission of schistosomiasis: an introductory view. *Am J Trop Med Hyg.* 1996;55(Suppl):135–43.
29. Yang Y, Xiao D. A mathematical model with delays for schistosomiasis japonicum transmission. *Chinese Ann Math.* 2010;31B:433–46.
30. Gao SJ, He YY, Liu YJ, Yang GJ, Zhou XN. Field transmission intensity of *Schistosoma japonicum* measured by basic reproduction ratio from modified Barbour's model. *Parasit Vectors.* 2013;6:141.
31. Wu KC. Mathematical model and transmission dynamics of schistosomiasis and its application. *Chinese Trop Med.* 2005;5:837–44.
32. Zhao J, Ouyang T, Ding G. Numerical evaluation of new control method for schistosomiasis. *Chinese J Biomed Eng.* 2011;28:725–31.
33. Chen XQ, Xu MG. Analysis of the infection in cattle schistosomiasis in Hubei. *Hubei J Anim Vet Sci.* 1993;1:23–4.
34. Mao SB. Schistosome biology and control of schistosomiasis. Beijing: People's Health Publication House; 1990.
35. Bacaër N, Guernaoui S. The epidemic threshold of vector-borne diseases with seasonality. *J Math Biol.* 2006;53:421–36.
36. Bacaër N. Approximation of the basic reproduction number  $R_0$  for vector-borne diseases with a periodic vector population. *Bull Math Biol.* 2007;69: 1067–91.
37. Wang WD, Zhao XQ. Threshold dynamics for compartmental epidemic models in periodic environments. *J Dyn Differ Equ.* 2008;20:699–717.
38. Zhou XN, Bergquist R, Leonardo L, Yang GJ, Yang K, Sudomo M, Olveda R. Schistosomiasis japonica control and research needs. *Adv Parasitol.* 2010;72: 145–78.
39. Liang S, Yang C, Zhong B, Qiu D. Re-emerging schistosomiasis in hilly and mountainous areas of Sichuan, China. *Bull World Health Organ.* 2006;84: 139–44.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

